

---

## Highly parallel genome-wide expression analysis of single mammalian cells.

**Journal:** PLoS One

**Publication Year:** 2012

**Authors:** Jian-Bing Fan, Jing Chen, Craig S April, Jeffrey S Fisher, Brandy Klotzle, Marina Bibikova, Fiona Kaper, Mostafa Ronaghi, Sten Linnarsson, Takayo Ota, Jeremy Chien, Louise C Laurent, Jeanne F Loring, Sean V Nisperos, Gina Y Chen, Jiang F Zhong

**PubMed link:** 22347404

**Funding Grants:** Ensuring the safety of cell therapy: a quality control pipeline for cell purification and validation

### Public Summary:

**BACKGROUND:** We have developed a high-throughput amplification method for generating robust gene expression profiles using single cell or low RNA inputs. **METHODOLOGY/PRINCIPAL FINDINGS:** The method uses tagged priming and template-switching, resulting in the incorporation of universal PCR priming sites at both ends of the synthesized cDNA for global PCR amplification. Coupled with a whole-genome gene expression microarray platform, we routinely obtain expression correlation values of  $R(2) \sim 0.76-0.80$  between individual cells and  $R(2) \sim 0.69$  between 50 pg total RNA replicates. Expression profiles generated from single cells or 50 pg total RNA correlate well with that generated with higher input (1 ng total RNA) ( $R(2) \sim 0.80$ ). Also, the assay is sufficiently sensitive to detect, in a single cell, approximately 63% of the number of genes detected with 1 ng input, with approximately 97% of the genes detected in the single-cell input also detected in the higher input. **CONCLUSIONS/SIGNIFICANCE:** In summary, our method facilitates whole-genome gene expression profiling in contexts where starting material is extremely limiting, particularly in areas such as the study of progenitor cells in early development and tumor stem cell biology.

### Scientific Abstract:

**BACKGROUND:** We have developed a high-throughput amplification method for generating robust gene expression profiles using single cell or low RNA inputs. **METHODOLOGY/PRINCIPAL FINDINGS:** The method uses tagged priming and template-switching, resulting in the incorporation of universal PCR priming sites at both ends of the synthesized cDNA for global PCR amplification. Coupled with a whole-genome gene expression microarray platform, we routinely obtain expression correlation values of  $R(2) \sim 0.76-0.80$  between individual cells and  $R(2) \sim 0.69$  between 50 pg total RNA replicates. Expression profiles generated from single cells or 50 pg total RNA correlate well with that generated with higher input (1 ng total RNA) ( $R(2) \sim 0.80$ ). Also, the assay is sufficiently sensitive to detect, in a single cell, approximately 63% of the number of genes detected with 1 ng input, with approximately 97% of the genes detected in the single-cell input also detected in the higher input. **CONCLUSIONS/SIGNIFICANCE:** In summary, our method facilitates whole-genome gene expression profiling in contexts where starting material is extremely limiting, particularly in areas such as the study of progenitor cells in early development and tumor stem cell biology.

---

**Source URL:** <https://www.cirm.ca.gov/about-cirm/publications/highly-parallel-genome-wide-expression-analysis-single-mammalian-cells>